

---

## Problem Set 2 Quantum statistics

---

### 1 Two-dimensional electron gas

A confined electron gas can form at the interface between two doped semiconductors (e.g., GaAs/AlGaAs). The confinement is such that one can consider that the gas is strictly two-dimensional. Electron-electron interactions will be neglected in the following and we will adopt the effective mass approximation. We call  $n$  the electronic density of the gas and  $A = L_x L_y$  its surface (which we assume to be very large as compared to all the other length scales of the problem). Here,  $L_x$  and  $L_y$  are the lateral dimensions of the gas in the  $x$  and  $y$  directions, respectively. We recall that the electrons are spin-1/2 fermions, and thus obey the Fermi-Dirac statistics. The average occupancy of an energy state  $\epsilon$  is then given by the Fermi-Dirac distribution function

$$f(\epsilon) = \frac{1}{e^{\beta(\epsilon-\mu)} + 1}, \quad (1)$$

where  $\beta = 1/k_B T$ , with  $T$  the temperature of the gas, and where  $\mu = \mu(T)$  is the chemical potential.

- (a) Plot the Fermi-Dirac distribution (1). In particular, analyze the  $T = 0$  case.
- (b) Using periodic boundary conditions (why can you do so?), solve Schrödinger's equation and show that the electronic dispersion is given by

$$\epsilon_{\mathbf{k}} = \frac{\hbar^2 |\mathbf{k}|^2}{2m},$$

where the wavevector  $\mathbf{k} = (k_x, k_y)$  is quantized according to  $k_x = 2\pi n_x/L_x$  and  $k_y = 2\pi n_y/L_y$ , with  $n_x$  and  $n_y$  integer numbers.

- (c) Show that the electronic density of states  $\rho(\epsilon)$  is energy-independent and is given by  $\rho(\epsilon) = 1/\Delta$ , where  $\Delta = \pi\hbar^2/mA$ .
- (d) Give an expression for the average number  $N$  of electrons in the gas. Deduce from the previous result that the chemical potential reads

$$\mu(T) = k_B T \ln \left( e^{T_F/T} - 1 \right),$$

where  $T_F$  is the Fermi temperature, defined through the Fermi energy as  $E_F = k_B T_F$ . What is the definition of the Fermi energy? Give an expression of  $E_F$  as a function of  $N$  and  $\Delta$ . Interpret this result. Plot  $\mu$  as a function of  $T$ .

- (e) Give a formal expression of the average energy  $E$  of the system in terms of an integral over  $\epsilon$ , that we will not explicitly calculate. Show that the grand-canonical potential reads

$$\Omega = -\frac{k_B T}{\Delta} \int_0^\infty d\epsilon \ln \left( 1 + e^{-\beta(\epsilon-\mu)} \right).$$

Deduce from the previous two results that  $\Omega = -E$ .

- (f) Show that the two-dimensional pressure  $P$  of the gas is related to the average energy via the expression  $P = E/A$ .
- (g) Using your answers to questions (e) and (f) above, derive the equation of state at  $T = 0$ .

- (h) At low temperature ( $T \ll T_F$ ), expand the average energy to second order in  $T/T_F$  so as to obtain the equation of state. Notice that

$$\int_{-\infty}^{+\infty} dx \frac{x^2 e^x}{(e^x + 1)^2} = \frac{\pi^2}{3}.$$

- (i) Calculate the equation of state at high temperature ( $T \gg T_F$ ). Comment your result.  
 (j) (*Optional question*) Calculate now the equation of state for an arbitrary temperature. One gives

$$\int_0^{\infty} dx \frac{x}{e^{x/a} + 1} = -\text{Li}_2(-a),$$

where  $a$  is a constant, and where  $\text{Li}_s(z) = \sum_{k=1}^{\infty} z^k/k^s$  is the polylogarithm function of order  $s$ .

- (k) Compare all the results of this problem to the three-dimensional case encountered in the lecture.

## 2 Bose–Einstein condensation

Let us consider a system of  $N$  bosons with mass  $m$  and spin  $s$  ( $s \in \mathbb{N}$ ) occupying a volume  $V$ . In the following, the interactions between the bosons are neglected. Accessible energy levels are denoted by  $\epsilon_{\mathbf{k}}$ , and the ground state energy is set to zero.

- (a) What is the average occupancy  $n(\epsilon)$  of a state of energy  $\epsilon$  at temperature  $T$ ? Show that the density of states takes the form  $d(\epsilon) = KV\sqrt{\epsilon}$ , where  $K$  is a constant. Give the expression for  $K$ . What is the sign of the chemical potential  $\mu$ ? How is  $\mu$  determined in the thermodynamic limit?  
 (b) Plot on the same graph  $n(\epsilon)$  as a function of  $\epsilon$  for two different chemical potentials  $\mu_1 < \mu_2$  while  $T$  is being kept fixed. On another graph, plot  $n(\epsilon)$  as a function of  $\epsilon$  for two different temperatures  $T_1 < T_2$  while  $\mu$  is being kept fixed. Considering that the number of particles is fixed, show that

$$\left( \frac{\partial \mu}{\partial T} \right)_N < 0.$$

- (c) By introducing the fugacity  $\varphi = e^{\beta\mu}$  as well as the function

$$f(\varphi) = \int_0^{\infty} dx \frac{\sqrt{x}}{e^{x/\varphi} - 1},$$

determine graphically the chemical potential  $\mu$ . What happens when the temperature is lowered? Show that there exists a critical temperature  $T_B$ , called the *Bose temperature*, for which  $\mu = 0$ . Note that

$$\int_0^{\infty} dx \frac{\sqrt{x}}{e^x - 1} = \frac{\sqrt{\pi}}{2} \zeta\left(\frac{3}{2}\right),$$

where  $\zeta(z)$  is the Riemann zeta function, which is defined for any complex number  $z$  such that  $\text{Re}(z) > 1$  by the Riemann series  $\zeta(z) = \sum_{n=1}^{\infty} n^{-z}$ . In particular,  $\zeta(3/2) \simeq 2.61$  and  $\zeta(5/2) \simeq 1.34$ .

- (d) We now consider that  $T < T_B$  and we assume  $N$  to be fixed. Show that the number of particles in the ground state is given by

$$N_0 = N \left[ 1 - \left( \frac{T}{T_B} \right)^{3/2} \right].$$

Is it possible to condensate photons?

- (e) Give an expression of the average energy  $E$  of the system in terms of an integral over  $\epsilon$  that we will not explicitly calculate. Show that the grand-canonical potential  $\Omega$  reads

$$\Omega = KVk_{\text{B}}T \int_0^\infty d\epsilon \sqrt{\epsilon} \ln \left( 1 - e^{-\beta(\epsilon-\mu)} \right).$$

Deduce from the two previous results that  $\Omega = -2E/3$ . Then, show that the pressure of the Bose gas is given by  $P = 2E/3V$ .

- (f) Derive an expression for the pressure of the system at  $T < T_{\text{B}}$ . Note that

$$\int_0^\infty dx \frac{x^{3/2}}{e^x - 1} = \frac{3\sqrt{\pi}}{4} \zeta \left( \frac{5}{2} \right).$$

- (g) We now consider that  $T$  is kept constant, instead of  $V$  ( $N$  remains fixed throughout). What happens when the volume of the system is decreased? Show that the Bose condensation takes place for

$$V_{\text{B}} = \frac{1}{(2s+1)\zeta(3/2)} N \Lambda_T^3,$$

where  $\Lambda_T = (2\pi\hbar^2/mk_{\text{B}}T)^{1/2}$  is the thermal de Broglie wavelength. Plot a few isothermal curves in a  $P$ - $V$  diagram. Discuss your results.

- (h) Liquid  ${}^4\text{He}$  presents a superfluid transition at 2.17 K. Compare such an experimental result to the Bose temperature. Parameters for liquid  ${}^4\text{He}$  are: spin  $s = 0$ , density  $0.12 \text{ g/cm}^3$ , and  $m = 4 \times m_{\text{proton}} = 6.7 \times 10^{-27} \text{ kg}$ . We recall that  $\hbar = 1.0 \times 10^{-34} \text{ J.s}$  and  $k_{\text{B}} = 1.4 \times 10^{-23} \text{ J/K}$ .